

A new species and notes on *Baccharis* sect. *Caulopterae* (Asteraceae) from Brazil

Uma nova espécie e notas sobre Baccharis L.
sect. Caulopterae DC. (Asteraceae) no Brasil

Angelo Alberto Schneider¹, Gustavo Heiden² & Ilisi Iob Boldrini¹

Abstract

A new species of *Baccharis* L. sect. *Caulopterae* DC. (Asteraceae) from sandy soils and dunes on the coastal plain of Rio Grande do Sul state, Brazil, is presented: *Baccharis dunensis* A.A.Schneid. & G.Heiden. The new species is described, illustrated and compared to more closely related species. Additionally, *B. burchelli* Baker is reinstated, new synonyms are proposed for *B. burchellii* and *B. crispa* Spreng., and a lectotype is designated for *B. regnellii* Sch.Bip. ex Baker.

Key words: Astereae, *Baccharis* subg. *Molina*, Compositae, Trimeria group.

Resumo

Uma nova espécie de *Baccharis* L. sect. *Caulopterae* DC. (Asteraceae) dos solos arenosos e dunas da planície costeira do litoral do Rio Grande do Sul, Brasil, é apresentada: *Baccharis dunensis* A.A. Schneid. & G. Heiden. A nova espécie é descrita, ilustrada e comparada com as espécies mais próximas. Adicionalmente, o binômio *B. burchelli* Baker é restabelecido, novos sinônimos são propostos para *B. burchellii* e *B. crispa* Spreng., e um lectótipo é designado para *B. regnellii* Sch.Bip. ex Baker.

Palavras-chave: Astereae, *Baccharis* subg. *Molina*, Compositae, grupo Trimeria.

Introduction

Baccharis is a new world genus comprising about 360 species (Nesom & Robinson 2007), distributed from southern Canada to southern South America (Fielding 2001; Giuliano 2001). The tufted indumentum of the leaves and stems, with adjoining basal cells of the trichomes, and the occurrence of dioecy are probably synapomorphic characters of *Baccharis* (Müller 2006).

Baccharis sect. *Caulopterae* DC. is represented by about 30 species restricted to South America (Barroso 1976; Giuliano 2001; Müller 2006) and is characterized mainly due to presence of winged stems, epaleaceous clinanthia densely covered with biseriate glandular hairs, papillose glabrous achenes, and pappus bristles of female flowers enlarged basally and fused into a ring.

During the preparation of the checklist of Brazilian species of *Baccharis* the description of a new species and the publication of some nomenclatural notes on *Baccharis* sect. *Caulopterae* DC. became necessary.

Results and Discussion

A new species of the coastal dunes of Rio Grande do Sul State, Southern Brazil

Baccharis dunensis A.A.Schneid. & G.Heiden, sp. nov. Type: BRASIL. RIO GRANDE DO SUL: Capão da Canoa, entre Noiva do Mar e Atlântida, 12.IV.1978, J. Pfadenhauer 250 (holotype ICN!). Fig. 1.

“*Baccharis riograndensis* Teodoro & J.E.Vidal” affinis, sed alis angustioribus et pappo pluriseriato differt, similis *B. triangularis*

¹Universidade Federal do Rio Grande do Sul, Programa de Pós-Graduação em Botânica, Av. Bento Gonçalves 9500, 91501-970, Porto Alegre, RS, Brasil.
angeloschneider@yahoo.com.br and ilsi.boldrini@ufrgs.br

²Universidade de São Paulo, Programa de Pós Graduação em Botânica, Depto. Botânica, Instituto de Biociências, R. do Matão, Tl. 14, 321,05508-900, São Paulo, SP, Brasil. gustavo.heiden@gmail.com




Figure 1 – *Baccharis dunensis* A.A.Schneid. & G. Heiden – a. habit; b. staminate capitulum; c. staminate flower (pappus removed); d. pistillate capitulum; e. phyllaries of pistillate capitulum; f. corolla and style of pistillate flower; g. cypselae. (a, d-g Schwacke 283; b, c Heiden 1054).

Hauman loco habitato, habitu, caule anguste alato, sed inflorescentia spiciformi et foliis apicalibus coronatis abscentibus differt.

Subshrub, 0.1–0.3 m tall; branching sympodial, shoots erect or erectopatente, ending in a capitulecence, green, resinous; stems 3-winged, wings to 0.1–2 mm wide, plane, green, attenuate near to the nodes, internodia 0.3–2 cm long. Leaves squamiform, sessile, nearly triangular, 0.1–0.7 × 0.1–0.7 mm. Capitula sessile, solitary, along terminal spike-like branches 1–4 cm long. Male capitula 5.5–7 mm long; flowers 10–25; involucre 5–6 × 2–4 mm, campanulate; phyllaries in 4 or 5 series, outermost phyllaries oblong, median phyllaries broadly lanceolate, innermost phyllaries lanceolate, all phyllaries with apex obtuse, margins yellow to light brown, broadly scarious; corolla 4–5 mm long, tube 2–3 mm long, lobes 1.5–2 mm long, coiled at maturity; style slightly exceeding the corolla, 4.5–6 mm long, the apex nearly fully divided into lanceolate branches, dilated towards apex, ovary abortive, glabrous and reduced; pappus uniseriate, 5–6 mm long, bristles 15–25, apically broadened and scarcely barbellate. Female capitula 7.5–12 mm long; flowers 10–30; involucre 7–8 × 1.5–2.5 mm, cylindrical; phyllaries in 4–6 series, like those of male capitula; corolla 5–6 mm long, filiform, short-ligulate, the ligule denticulate, glabrous; style 6.5–7.3 mm long; cypselae 1.5–1.7 × 0.2–0.4 mm, nearly cylindric, epapillate; 10–12-ribbed, pappus pluriseriate to several seriate, 3.5–8 mm long, bristles 37–60, persistent, basally fused.

Material examined: RIO GRANDE DO SUL: Mostardas, Lagoa do Peixe, 21.II.1970, ♂, E. Viana et al. s.n. (ICN 7531). Rio Grande, Ilha dos Marinheiros, 29.II.1880, ♀, C.A.W. Schwacke 283(R); 25.V.2009, ♂, G. Heiden et al. 1053 (HECT, ICN, JE, K, RB). São José do Norte, I.1992, ♂, P. Tagliani 77 (HURG); 10.IX.2008, G.H. Silveira & P.C. Crespan s.n. (ICN 157830). Tavares, Lagoa do Peixe, 15.XII.1986, ♀, E. Danilevitz 38 (HAS). Tramandaí, 20.VIII.2005, B. Irgang s.n. (ICN 157829).

Baccharis dunensis is a small subshrub (0.1–0.3 m tall), with branched and narrow winged alate stems. It is likely to *B. triangularis* because both have similar habit and habitat and narrow winged stems. However, it can be distinguished from the later species mainly due the absence of a pseudoinvolucre of reduced bractlike cuneate leaves, crowded around the capitula, which are apically solitary on the fertile branches. It is also

similar to *B. riograndensis* by the cylindrical-elongated female capitula, but it is distinguished by the smaller tall (0.1–0.3 m to *B. dunensis* and 0.3–0.6 m to *B. riograndensis*), the narrow wings of the vegetative branches (0.1–2 vs. 2.5–7 mm wide to *B. riograndensis*) and pappus several seriate.

Baccharis dunensis is probably endemic to the coastal plain of the state of Rio Grande do Sul. It is restricted to sandy soils and dunes along the coastal plain. Flowering and fruiting specimens have been collected from January to May. The conservation status was framed in VU Ba(iii)– Vulnerable, according to *IUCN Red list category* (IUCN 2009) because the species is known to less than 10 locations and these places are suffering intense anthropogenic pressure resulting in loss of area and decline of habitat quality, mainly due to urban expansion, dunes removal and biological invasion of coastal vegetation by exotic species such as *Casuarina* L. and *Pinus* L.

Nomenclatural notes on *Baccharis* sect. *Caulopterae* from Brazil

Baccharis burchellii Baker in Martius, Eichler & Urban, Fl. bras. 6(3): 44. 1882. Type: BRAZIL. SÃO PAULO: "ad marginem silvulae prope Morumbi", W.J. Burchell 4471 (holotype K, photo ICN!; isotype P, photo 1CN!).

Baccharis regnellii var. *subalata* Heering in Usteri, Fl. São Paulo 258. 1911. Type: BRAZIL. SÃO PAULO: 23.XI.1906, A. Usteri 70b (holotype not found; isotype SP!), *syn. nov.*

Baker (1882) assigned *Baccharis burchellii* (ser. *Caulopterae*) and *B. regnellii* (ser. *Oblongifoliae*) in distinct series, based on the presence/absence of winged stems and leaf shape. Barroso (1976) recognized the affinities between these species, however she considered *B. burchellii* as a synonym of *B. regnellii* var. *subalata* Heering. In the current work, *B. burchellii* was considered as a distinct species and its name has been applied based on their studied types and recent collections. *Baccharis burchellii* is morphologically related to *B. regnellii*, but it can be distinguished mainly by the winged stem and sessile leaves with entire margins (vs. stem apterous and leaves with petiole and crenulate margins in *B. regnellii*).

Baccharis crispa Spreng., Syst. veg. 3: 466. 1826.
Molina crispa (Spreng.) Less., Linnaea 6: 141. 1831.
Baccharis genistelloides var. *crispa* (Spreng.) Baker in Martius, Eichler & Urban, Fl. bras. 6(3): 41. 1882. *Pingraea crispa* (Spreng.) F.H. Hellw., Candollea 46: 217. 1993. *Baccharis genistelloides* subsp. *crispa* (Spreng.) Joch. Müll., Syst. Bot. Monogr. 76: 198. 2006. Type: Uruguay, Montevideo, 1821-1822, C, F. Sellow d397 [M.I.B. 736] (holotype P, photo ICN!; isotypes BR; G; G-DC; R!, W).

Baccharis subcrispa Teodoro, Contr. Inst. Geobiol. 8: 39. 1957. Type: BRAZIL. RIO DE JANEIRO: Pedra do Altar, A.C. Brade 15593 (holotype RB!; isotypes ICN!, JE!), syn. nov.

Barroso (1976) considered *Baccharis subcrispa* as synonym of *B. opuntioides* Mart. ex Baker, an endemic species to the Serra do Caparaó (ES/MG). However, based on the analysis of the holotype of *B. subcrispa*, we ascertain this name as conspecific to *B. crispa* due its morphological congruence.

Baccharis regnelli Sch.Bip. ex Baker in Martius, Eichler & Urban, Fl. bras. 6(3): 74. 1882. Type: BRAZIL. MINAS GERAIS: Poços de Caldas, 9.IX.1867, ♀, A.F. Regnell II-155. (lectotype R!, here designated; duplicates BR, photo ICN!; C, photo ICN!; US, photo ICN!).

Baker (1882) described *Baccharis regnelli* listing two specimens (*Regnell II.155* and *Glaziovii 7718*). During the revision of herbarium R, a duplicate of the gathering *Regnell II.155* (R) was found. This specimen is designated here as lectotype, due its congruency with the original description and by representing a fertile branch with developed capitula and flowers.

Acknowledgements

The authors are grateful to the herbaria curators, to the collectors of the studied specimens, to CAPES and CNPq-PROTAX for the financial support of our studies, to João Iganci for preparing the illustration, to Tarçiso Filgueiras for revision of the Latin diagnosis and to the anonymous reviewers for contributions on the improvement of the text.

References

- Baker, J.G. 1882. Compositae. III. Asteroideae, Inuloideae. In: Martius, C.F.P. von; Eichler, A. W. & Urban, I. (ed.). *Flora brasiliensis*. München, Wien, Leipzig. Vol. 6. Pp. 1-442.
- Barroso, G.M. 1976. Compositae, subtribo Baccharidinae Hoffmann. Estudo das espécies ocorrentes no Brasil. Rodriguésia 28: 3-273.
- Fielding, R.R. 2001. *Baccharis*: a genus of the Asteraceae new to Canada. Proceedings of the Nova Scotian Institute of Science 41: 214-215.
- Giuliano, D.A. 2001. Clasificación infragenérica de las especies argentinas de *Baccharis* (Asteraceae, Astereae). Darwiniana 39: 131-154.
- IUCN. 2009. IUCN Red list of threatened species. Version 2009.1. Available in <<http://www.iucnredlist.org>>. Access on 15 November 2009.
- Müller, J. 2006. Systematics of *Baccharis* (Compositae-Astereae) in Bolivia, including an overview of the genus. Systematic Botany Monographs 76: 1-341.
- Nesom, G. & Robinson, H. [2006] 2007. XV. Tribe Astereae Cass. In: Kadereit, J. W. & Jeffrey, C. (eds.). The families and genera of vascular plants. Kubitzki, K. (series ed.). Flowering plants: Eudicots: Asterales. Vol. VIII. Springer, Berlin. Pp. 284-342.